

Strong CP θ -problem in QCD and local gluon condensate

V.V.Kiselev*

*Russian State Research Center "Institute for High Energy Physics",
Protvino, Moscow Region, 142281, Russia
Fax: +7-0967-744739*

We calculate the contribution of local gluon condensate into the quadratic θ -term of vacuum energy and discuss its application to the study of axion mass.

PACS numbers: 12.38.Aw, 14.80.Mz, 11.30.Er

I. INTRODUCTION

The strong CP problem reveals itself by the term of lagrangian permitting the CP-violation in an explicit form

$$\mathcal{L}_\theta = -\theta \frac{\alpha_s}{8\pi} \epsilon_{\mu\nu\alpha\beta} G_a^{\mu\nu}(x) G_a^{\alpha\beta}(x), \quad (1)$$

where *a priori* free quantity θ denotes the effective parameter including the bare contribution as well as the term caused by the matrix of quark masses M ,

$$\theta = \theta_{bare} - \arg(\det M).$$

The smallness of θ following from the experimental observations, basically, by the constraints on the electric dipole moment of neutron, looks rather accidental in QCD, unless we do not involve some additional mechanisms. The most elegant opportunity is the Peccei–Quinn symmetry [1] introducing an axion field coupled to the CP-odd product of gluon strength tensors in (1). In that case, the quantity θ gains the dynamical nature due to the axion a , so that $\theta \rightarrow \theta + a/f_a$, where we have introduced the axion decay constant f_a , and the problem is solved by an appropriate effective potential for θ , if the minimal energy is reached at $\langle\theta\rangle = 0$. Since the bare and quark mass terms in θ lead to a global additive redefinition of axion field, we can easily see that the positive mass squared of axion in the effective action is a necessary condition in order to control the stability, while the position of stable point should be at $\theta = 0$ to get the above solution of strong CP-problem.

The real value of axion mass depends on whether the axion is coupled to the quark fields or not. The corresponding term of fermion contribution is given by

$$\mathcal{L}_f = i g_n a \bar{q}_n \gamma_5 q_n, \quad (2)$$

where n marks the quark flavor [2]. There is a possibility to introduce the coupling to the leptons, too, [3]. So, due to the fermion component the mixing of axion with the pseudo-Goldstone bosons appearing under the spontaneous breaking of chiral symmetry results in the

following general formula [4] for the axion mass:

$$m_a^2 = \frac{N_a^2}{f_a^2} \frac{\langle \bar{q}q \rangle K}{\langle \bar{q}q \rangle - K \text{Tr} M^{-1}}, \quad (3)$$

where N_a is the color anomaly of Peccei–Quinn current, which we put equal to 1, $\langle \bar{q}q \rangle$ denotes the quark condensate¹, while K is a dimensional quantity ordinary represented by the instanton-induced term in the effective potential. Therefore, the light quarks coupled to the axion essentially change the situation, since they suppress purely gluon contribution. Nevertheless, the consideration of gluon term in the quadratic θ -action remains actual as the leading correction to the quark condensate, if the axion is coupled to the quarks, or as the main term, if the axion is coupled to the gluons, only.

In the present study we calculate the effective action quadratic in θ due to the contribution of the local gluon condensate. This contribution can be competitive with the term induced by the instantons, since, first, as observed by the lattice simulations, the instantons do not dominate in the determination of vacuum properties in QCD because the vacuum is too warm, second, modelling the gluon condensate by the dilute instanton gas is problematic as the gas should consist of rather wide and dense instantons in order to reach the observed magnitude of the gluon condensate. In addition, the distribution over the instanton sizes involves extra model-parameters. Thus, the gluon condensate contribution itself is of interest.

II. CALCULATIONS AND RESULTS

In the calculation of diagram shown in Fig. 1 we take the leading term caused by the local gluon condensate,

$$\begin{aligned} \langle G_a^{\mu\nu}(x) G_b^{\mu'\nu'}(y) \rangle &= \langle G_a^{\mu\nu}(0) G_b^{\mu'\nu'}(0) \rangle \\ &+ y^\alpha \langle G_a^{\mu\nu}(0) \partial_\alpha G_b^{\mu'\nu'}(0) \rangle + x^\alpha \langle \partial_\alpha G_a^{\mu\nu}(0) G_b^{\mu'\nu'}(0) \rangle \\ &+ x^\alpha y^\beta \langle \partial_\alpha G_a^{\mu\nu}(0) \partial_\beta G_b^{\mu'\nu'}(0) \rangle + \dots \\ &= \langle G_a^{\mu\nu}(0) G_b^{\mu'\nu'}(0) \rangle + \text{higher condensates} \end{aligned} \quad (4)$$

*Electronic address: kiselev@th1.ihep.su

¹ Further, one usually insets the relation $(m_u + m_d) \langle \bar{q}q \rangle = -m_\pi^2 f_\pi^2$ with $\langle \bar{q}q \rangle = \langle \bar{u}u \rangle + \langle \bar{d}d \rangle$ and $f_\pi \approx 93.3$ MeV.

so that we can write down its contribution to the effective action in the explicit form

$$i\Gamma_{\theta\theta} = \lim_{k \rightarrow 0} (-i)^2 \left(\frac{\alpha_s}{8\pi} \right)^2 \epsilon_{\mu\nu\alpha\beta} \epsilon_{\mu'\nu'\alpha'\beta'} \langle G_a^{\mu\nu} G_b^{\mu'\nu'} \rangle \times \\ ik^\alpha (-ik^{\alpha'}) \frac{-ig^{\beta\beta'}}{k^2} \delta_{ab} F_{comb}, \quad (5)$$

where the combinatoric factor $F_{comb} = 16$ is determined by the permutations of gluon strength tensors and the evident relation $\epsilon_{\mu\nu\alpha\beta}(\partial^\alpha A_a^\beta - \partial^\beta A_a^\alpha) = 2\epsilon_{\mu\nu\alpha\beta}\partial^\alpha A_a^\beta$.




FIG. 1: The quadratic θ -term caused by the local gluon condensate.

For the vacuum expectation value we evidently have the following general expression conserving the color and Lorentz symmetries:

$$\langle G_a^{\mu\nu} G_b^{\mu'\nu'} \rangle = \frac{\delta_{ab}}{96} (g^{\mu\mu'} g^{\nu\nu'} - g^{\mu\nu'} g^{\nu\mu'}) \langle G_d^{\mu\nu} G_{\mu\nu}^d \rangle. \quad (6)$$

Then, we find the gluon condensate part of quadratic term in the effective potential for the global phase θ

$$V_{\theta^2}[G^2] = \theta^2 \left\langle \frac{\alpha_s^2}{8\pi^2} G_a^{\mu\nu} G_{\mu\nu}^a \right\rangle, \quad (7)$$

while the linear term due to the bare lagrangian in (1) does not contribute into the vacuum energy, since the physical vacuum infers condition (6).

We can straightforwardly get the contribution given by the local quark condensate into the quadratic term of axion in the way analogous to the described calculation with the gluon condensate, so that

$$V_{a^2}[\bar{q}q] = -a^2 \sum_n \frac{|g_n|^2}{m_n} \langle \bar{q}_n q_n \rangle, \quad (8)$$

which can be further transformed after the substitution for the universal coupling of the axion to the quarks [4]

$$g_n = \frac{1}{f_a} \frac{1}{\text{Tr } M^{-1}}, \quad (9)$$

expressing a consistency for the definition of axion decay constant through the calculation of axial anomaly and the absence of mass mixing between the axion and neutral pseudo-Goldstone mesons π^0 and η . Here, M denotes

the diagonal mass matrix of light quarks (u, d). Then, we get

$$V_{a^2}[\bar{q}q] = -\frac{1}{2} \frac{a^2}{f_a^2} \langle \bar{q}q \rangle \frac{m_u m_d}{m_u + m_d}, \quad (10)$$

which reproduces the result of [4] in the case of quark condensate dominance. We stress that the mixing of the axion with the flavor-singlet meson leads to the quadratic axion potential tending to zero in the limit of massless light quark in agreement with the general formula given by (3) despite the calculated term due to the local gluon condensate.

Indeed, comparing (10) with (7) after the substitution of $\theta \rightarrow a/f_a$ we could observe that the contribution by the local gluon condensate dominates in the axion mass. The picture is critically changed, if we take into account the neutral singlet of SU(3) flavor group. This singlet is coupled to the gluon anomaly, too, in the same manner as the axion with an analogous constant f_s instead of f_a , which produces the mixing of masses for the axion and singlet [4]. The resulting effect has been described in the Introduction, and we return to the quark condensate dominance in the axion mass up to small corrections. The description of above mechanism is given in [4] in detail.

Nevertheless, we stress that expression (7) gives explicit result in contrast to rather indefinite instanton-induced potential.

Let us discuss the renormalization group (RG) properties of results obtained. First, we can easily see that the contribution by the quark condensate (10) enters into the axion mass in the RG invariant form, since the ratio of quark masses and the product of quark mass with the condensate are invariant under the action of RG. Second, the gluon condensate term due to (7) reveals the RG dependence with nonzero anomalous dimension in QCD. However, this anomalous dimension is in the next α_s -order under study. Therefore, we have no any contradiction with the QCD principles in the given accuracy over α_s .

We do not make some numerical estimates, since they would be rather qualitative under assumptions on the effective value of α_s .

Thus, we clarify the physical meaning of our calculations and results.

III. CONCLUSION

We have calculated the contribution by the local gluon condensate into the effective potential quadratic over the θ determining the CP-violating term of the gluon anomaly in QCD. Such the effective potential gives the contribution into the axion mass providing the solution of strong CP-problem in the Peccei–Quinn mechanism. We expect the dominance of contribution caused by the local gluon condensate in comparison with the instanton-induced potential, which could be tested by the lattice simulations.

The axion plays an essential role in the connection of particle physics with the cosmology, in particular, as a dark matter candidate [5]. This fact makes the studies on its mass to be rather significant. We see that the QCD dynamics in such the investigations is of importance. We would like to point to some recent inter-

esting papers devoted to the problem under discussion: solutions of strong CP-problem [6, 7], on the Yang–Mills vacuum [8].

This work is in part supported by the Russian Foundation for Basic Research, grants 01-02-99315, 01-02-16585.

-
- [1] R. D. Peccei and H. R. Quinn, *Phys. Rev. D* **16**, 1791 (1977).
 - [2] J. E. Kim, *Phys. Rev. Lett.* **43**, 103 (1979);
M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Nucl. Phys. B* **166**, 493 (1980).
 - [3] M. Dine, W. Fischler and M. Srednicki, *Phys. Lett. B* **104**, 199 (1981);
A. R. Zhitnitsky, *Sov. J. Nucl. Phys.* **31**, 260 (1980) [*Yad. Fiz.* **31**, 497 (1980)].
 - [4] H. Georgi, D. B. Kaplan and L. Randall, *Phys. Lett. B* **169**, 73 (1986);
K. Choi, K. Kang and J. E. Kim, *Phys. Lett. B* **181**, 145 (1986).
 - [5] S. Khalil and C. Munoz, *Contemp. Phys.* **43**, 51 (2002) [arXiv:hep-ph/0110122].
 - [6] S. L. Glashow, arXiv:hep-ph/0110178.
 - [7] Z. Berezhiani, L. Gianfagna and M. Giannotti, *Phys. Lett. B* **500**, 286 (2001) [arXiv:hep-ph/0009290].
 - [8] V. Chernyak, arXiv:hep-th/9808092.